

# REGULATED CHIRAL GAUGE THEORIES.<sup>1</sup>

H. NEUBERGER

*Department of Physics and Astronomy, Rutgers University, Piscataway,  
NJ 08855, USA*

*E-mail: neuberg@physics.rutgers.edu*

Noncompact chiral abelian gauge theories are defined on the lattice using the overlap formalism. The main definitions are presented, the role of anomaly cancellation is discussed, and the triviality issue in four dimensions is explained.

## 1 Introduction

During the last several years we have learned how to put chiral gauge theories on the lattice. For recent reviews and further reference see [1]. The best understood case is abelian and aims at defining the theory on infinite Euclidean space  $R^d$  [2].

We start by defining the Lagrangian. We work in even  $d$ -dimensional Euclidean space, on an infinite hypercubic lattice. The action has a pure gauge part and a fermionic part. We shall start with the bosonic part, and next turn to the fermionic component. This shall be followed by a display of the lattice Wess Zumino action and of lattice anomalies. After that, trivial co-cycle components will be subtracted from the Wess Zumino action leading to the non-perturbative restoration of gauge invariance when anomalies cancel in perturbation theory. We then address the particular case of  $d = 4$  where anomalous and non-anomalous theories are both trivial, but in different ways. We conclude with a brief summary.

## 2 Photons

The pure gauge term is given by:

$$S_{\text{pure gauge}} = \frac{1}{4e^2} \sum_x F_{\mu\nu}^2(x) + \frac{m_\gamma^2}{2} \sum_x A_\mu^2(x) + \frac{1}{2\xi} \sum_x [\nabla_\mu^{x-} A_\mu(x)]^2. \quad (1)$$

The field strength is defined in terms of the noncompact vector potential  $A_\mu(x)$  by:

$$F_{\mu\nu}(x) = \nabla_\mu^{x+} A_\nu(x) - \nabla_\nu^{x+} A_\mu(x), \quad (2)$$

---

<sup>1</sup> INVITED TALK AT THE INTERNATIONAL CONFERENCE "QUANTIZATION, GAUGE THEORY, AND STRINGS, DEDICATED TO THE MEMORY OF PROFESSOR EFIM FRADKIN", JUNE 5 - JUNE 10, MOSCOW, RUSSIA.

where  $\nabla_\mu^{x+}$  are forward finite difference operators (the backward difference operators are  $\nabla_\mu^{x-} = (\nabla_\mu^{x+})^\dagger$ ). In addition, gauge configurations have vanishing probability in the path integral unless [3] for each site  $x$  and each Lorentz index pair  $\mu\nu$ ,

$$|F_{\mu\nu}(x)| \leq \eta, \quad (3)$$

where,

$$0 < \eta < \frac{1 - (1 + m)^2}{(1 + \frac{\sqrt{2}}{2})d(d - 1)}. \quad (4)$$

$m$  is a lattice parameter restricted by  $|1 + m| < 1$  (for our purposes here we can assume that  $m$  has been fixed at  $-1$ ). Although we have added a photon mass term and a gauge fixing term, gauge invariance plays a crucial role in the quantization of the theory, the consistency of its physical sector and the existence of a zero photon mass limit. So long  $m_\gamma^2 > 0$ ,  $A_\mu(x)$  goes to zero as  $|x| \rightarrow \infty$  for any gauge field configuration of non-zero probability. The entire gauge invariant content of the configuration is contained in the variables  $F_{\mu\nu}(x)$ . On a finite lattice with translationally invariant boundary conditions extra gauge invariant variables enter. These variables cause complications that we avoid here. So long  $0 < \xi < \infty$  the gaussian path integral over  $A_\mu(x)$  can be defined even before the restriction of equation (3) is enforced.

### 3 Chiral Fermions

The theory contains several Weyl fermions, either left or right handed, with charges  $q_1, q_2, \dots, q_n, 1$ ,  $|q_i| \leq 1$ ; at least one of the fermions has  $|q| = 1$ . For each fermion we define unitary matrices,

$$U_\mu(x; q) = e^{iqA_\mu(x)}, \quad (5)$$

and associated parallel transporters,

$$T_\mu(A)(\psi)(x) = U_\mu(x)\psi(x + \hat{\mu}) \quad (6)$$

acting on fermions  $\psi$ . The Wilson Dirac operator is given by:

$$D_W(A) = m + \sum_\mu (1 - V_\mu); \quad V_\mu = \frac{1 - \gamma_\mu}{2} T_\mu + \frac{1 + \gamma_\mu}{2} T_\mu^\dagger; \quad V_\mu^{-1} = V_\mu^\dagger \quad (7)$$

Associated to it, there is a hermitian Wilson Dirac operator

$$H_W(A) = \gamma_{d+1} D_W(A). \quad (8)$$

Here,  $\gamma_{d+1}$  generalizes the Hermitian  $\gamma_5$  matrix in four dimensions.

It is true then that the following unitary-hermitian operator [4] is well defined for all gauge configurations that can occur with non-zero probability:

$$\epsilon(A) = \text{sign}(H_W(A)). \quad (9)$$

To add the fermions we introduce an auxiliary Fock space [5] generated by

$$\{\hat{a}_\alpha^\dagger(x), \hat{a}_\beta(y)\} = \delta_{\alpha\beta} \delta_{xy}. \quad (10)$$

In this Fock space we define a Hamiltonian:

$$\hat{H}(A) = \hat{a}^\dagger \epsilon(A) \hat{a} + N_v, \quad (11)$$

where the constant  $N_v$  assures that the ground state energy of  $\hat{H}(A)$  be zero for all  $A$ .<sup>2</sup> There are local charge operators given by:

$$\hat{n}(x) = \frac{1}{2} [\hat{a}^\dagger(x) \hat{a}(x) - \hat{a}(x) \hat{a}^\dagger(x)]. \quad (12)$$

The total charge operator  $\hat{N} = \sum_x \hat{n}(x)$  commutes with the Hamiltonian. The ground state of  $\hat{H}(A)$  is unique and of zero charge for all  $A$ :

$$\hat{H}(A)|v(A)\rangle = 0, \quad \hat{N}|v(A)\rangle = 0. \quad (13)$$

Gauge transformations are generated by the exponentiation of the local charge operators. The resulting unitary operators are given by:

$$\hat{G}(\alpha) = e^{i \sum_x \alpha(x) \hat{n}(x)}. \quad (14)$$

The Hamiltonian transforms covariantly under a gauge transformation:

$$\hat{G}(\alpha) \hat{H}(A) \hat{G}^\dagger(\alpha) = \hat{H}(A^{(\alpha)}). \quad (15)$$

Here,  $A_\mu^{(\alpha)}(x) \equiv A_\mu(x) + \nabla_\mu^{x+} \alpha(x) \equiv A_\mu(x) + \alpha(x + \mu) - \alpha(x)$ , is a gauge transform of  $A_\mu(x)$ .

We also define a reference system, with  $\epsilon(A) \rightarrow \gamma_{d+1}$ ,  $\hat{H}(A) \rightarrow \hat{H}'$  and  $|v(A)\rangle \rightarrow |v'\rangle$ . Trivially,  $\hat{G}(\alpha)|v'\rangle = |v'\rangle$ . The phase of  $|v'\rangle$  is fixed in an arbitrary way. The phase of  $|v(A)\rangle$  is fixed by an adiabatic phase choice:

$$\langle v(tA) | \frac{dv(tA)}{dt} \rangle = 0; \quad \langle v' | v(0) \rangle > 0. \quad (16)$$

---

<sup>2</sup>Although the ground state energy of  $\hat{a}^\dagger \epsilon(A) \hat{a}$  is formally infinite because of the infinite volume, it is a meaningful fact that this energy is independent of  $A$ , for any  $A$ -background of non-zero probability. Moreover, all other eigenstates of  $\hat{H}(A)$  have trivial, integral, eigenvalues, which also do not depend on  $A$ .

The overlap provides a finite formula for the fermionic determinant:

$$\langle v' | v(A) \rangle. \quad (17)$$

This formula can be motivated by considering a system containing an infinite number of Dirac fermions with a mass matrix of the order of the ultraviolet cutoff, but with unit index. This interpretation connects the formula to the basic papers of Kaplan [6] and of Frolov and Slavnov [7].

## 4 Wess-Zumino action

If we made no phase choices, the overlap could be viewed as a line bundle over the space of gauge configurations. Because of the gauge transformation properties of the two states, this bundle descends into a line bundle over the space of gauge orbits. The phase choice can be viewed as a choice of section in the former, but, generically, not in the latter because the section may vary along gauge orbits [8]. Lack of ordinary gauge invariance is measured by the Wess-Zumino action  $\Phi(\alpha, A)$  which is given by

$$\langle v' | v(A^{(\alpha)}) \rangle = e^{i\Phi(\alpha, A)} \langle v' | v(A) \rangle. \quad (18)$$

Starting from the adiabatic phase choice, the Wess Zumino term can be explicitly calculated:

$$\Phi(\alpha, A) = - \sum_x \alpha(x) \int_0^1 dt \langle v(tA) | \hat{n}(x) | v(tA) \rangle. \quad (19)$$

$\Phi(\alpha, A)$  is linear in  $\alpha$  and vanishes for all  $\alpha$  if  $A = 0$ . The factor multiplying  $\alpha(x)$  is the local charge density and sums over the lattice to zero, for any  $A$ -configuration that occurs with non-zero probability.<sup>3</sup>

---

<sup>3</sup>It is amusing to note here that in earlier attempts to put chiral fermions on the lattice it was assumed that the decoupling of  $\alpha(x)$  on the trivial orbit was the central problem one was facing on the lattice. We now see that the lattice is not different from the continuum in this respect and whether the theory is anomalous or not, the gauge degrees of freedom can be made to decouple on the trivial orbit. This has no bearing on the existence of a continuum limit, which does depend on anomaly cancelation. Thus, as one should have known from the continuum, the main obstacles to the quantization of a chiral gauge theory on the lattice cannot be even identified before fermion loops are taken into account. However, the majority of previous attempts to put chiral fermions on the lattice used various forms of Yukawa type interactions and were deemed unsuccessful even before any fermion loop was taken into account.

## 5 Anomalies

The Bose symmetrized (“consistent”) anomaly comes directly from  $\Phi$ :

$$\Delta_{\text{cons}}(x; A) = - \int_0^1 dt \langle v(tA) | \hat{n}(x) | v(tA) \rangle. \quad (20)$$

One easily establishes that

$$\Delta_{\text{cons}}(x; A) = -\nabla_\mu^{x-} \mathcal{J}_{\mu \text{ cons}}(x; A); \quad \mathcal{J}_{\mu \text{ cons}}(x; A) = \frac{\partial \log \langle v' | v(A) \rangle}{\partial A_\mu(x)}. \quad (21)$$

In addition to the consistent current we have the covariant current [8]

$$\mathcal{J}_{\mu \text{ cov}}(x; A) = \frac{1}{\langle v' | v(A) \rangle} \langle v' | \frac{\partial v(A)}{\partial A_\mu(x)} \rangle_\perp, \quad (22)$$

with a covariant anomaly:

$$\Delta_{\text{cov}}(x; A) = -\nabla_\mu^{x-} \mathcal{J}_{\mu \text{ cov}}(x; A) = \langle v(A) | \hat{n}(x) | v(A) \rangle. \quad (23)$$

These currents are related to a current operator in Fock space,

$$[\hat{n}(x), \hat{H}(A)] = i \nabla_\mu^{x-} \hat{J}_\mu(x; A), \quad (24)$$

which admits the form:

$$\hat{J}_\mu(x; A) = - \frac{\partial \hat{H}(A)}{\partial A_\mu(x)}. \quad (25)$$

The relation to the c-number currents above is:

$$\langle v(A) | \hat{n}(x) | v(A) \rangle = \nabla_\mu^{x-} j_\mu(x; A), \quad (26)$$

where

$$j_\mu(x; A) = \int_0^1 dt \frac{i}{2} \left[ \langle v(tA) | \hat{J}_\mu(x; tA) | \frac{dv(tA)}{dt} \rangle - c.c. \right]. \quad (27)$$

$j_\mu(x; A)$  is not gauge invariant:

$$j_\mu(x; A^{(\alpha)}) - j_\mu(x; A) = \quad (28)$$

$$\frac{i}{2} \sum_y \left[ \int_0^1 dt \langle v(tA) | [\hat{J}_\mu(x; tA), \hat{J}_\nu(y; tA)] | v(tA) \rangle \right] \nabla_\nu^{y+} \alpha(y) \quad (29)$$

This introduces the Schwinger term [2]:

$$S_{\mu\nu}(x, y; A) = \langle v(A) | [\hat{J}_\mu(x; A), \hat{J}_\nu(y; A)] | v(A) \rangle. \quad (30)$$

Only the divergence of the Schwinger term enters in equation (29). The matrix element of the current commutator turns out to be Berry's curvature associated with the line bundle  $|v(A)\rangle$ , where  $A$  is viewed as "parameter space". (Note that this line bundle does not descend to a bundle over gauge orbits, but, its Berry curvature can be viewed as a two form over the space of gauge orbits.) Let us present the explicit expressions. The Berry connection is defined by

$$\mathcal{A}_{\mu x}(A) = \langle v(A) | \frac{\partial v(A)}{\partial A_\mu(x)} \rangle, \quad (31)$$

from which we get Berry's curvature [8]:

$$\mathcal{F}_{\mu x, \nu y}(A) = \frac{\partial \mathcal{A}_{\mu x}(A)}{\partial A_\nu(y)} - \frac{\partial \mathcal{A}_{\nu y}(A)}{\partial A_\mu(x)} = \quad (32)$$

$$\langle \frac{\partial v(A)}{\partial A_\mu(x)} | \frac{\partial v(A)}{\partial A_\nu(y)} \rangle - \langle \frac{\partial v(A)}{\partial A_\nu(y)} | \frac{\partial v(A)}{\partial A_\mu(x)} \rangle. \quad (33)$$

One can easily show that [2]:

$$S_{\mu\nu}(x, y; A) = 4\mathcal{F}_{\mu x, \nu y}(A). \quad (34)$$

The c-number covariant current can also be related to a matrix element of the current operator:

$$\mathcal{J}_{\mu \text{ cov}}(x; A) = -\frac{1}{2} \frac{\langle v' | \hat{J}_\mu(x; A) | v(A) \rangle}{\langle v' | v(A) \rangle}. \quad (35)$$

Both the covariant and consistent currents involve, in addition to  $|v(A)\rangle$ , also the reference state and are non-local functionals of  $A$ . The difference between the currents, given by Berry's curvature, does not involve  $|v'\rangle$ , and is a local functional of  $A$ .

## 6 Trivial cocycles

One can establish, by explicit construction or by more indirect means, the following fact: If the continuum anomaly cancelation condition is met, the

lattice Wess-Zumino functional is a locally trivial cocycle. To see what this means let us focus on two dimensions as an example [2]. Define a constant  $b$ :

$$b = \frac{1}{2} \epsilon_{\mu\nu} \sum_x S_{\mu\nu}(x, y; 0). \quad (36)$$

One can prove that there exist local gauge field functionals  $\chi_\rho$ , such that the divergence of the Schwinger term can be written as:

$$\nabla_\nu^{y-} S_{\mu\nu}(x, y; A) = \epsilon_{\mu\sigma} \nabla_\sigma^{x-} [\nabla_\rho^{y-} \chi_\rho(x, y; A) - b \delta_{x,y}]. \quad (37)$$

This reflects the identity  $\nabla_\mu^{x-} \nabla_\nu^{y-} S_{\mu\nu}(x, y; A) = 0$  which holds as a result of equations (24) and (30). Using  $\chi_\rho$ , the adiabatic phase choice of  $|v(A)\rangle$  can be augmented by adding to the adiabaticity requirement an explicit phase factor given by the exponent of a local, non-gauge invariant, functional of  $A_\mu(x)$

$$|v(A)\rangle \rightarrow e^{i\phi(A)} |v(A)\rangle, \quad (38)$$

which changes the Wess Zumino functional by a trivial cocycle,  $\Phi(\alpha, A) \rightarrow \Phi(\alpha, A) + \phi(A^{(\alpha)}) - \phi(A)$ , so that the new Wess Zumino functional becomes:

$$\Phi^{\text{new}}(\alpha, A) = -\frac{b}{2} \sum_x F_{12}(x) \alpha(x). \quad (39)$$

For the opposite handedness the overall sign gets reversed. It is now obvious that when the continuum anomaly cancelation condition  $\sum q_R^2 = \sum q_L^2$  is met the new Wess Zumino functional vanishes, showing that the previous Wess Zumino functional was a trivial cocycle. So, when anomalies cancel in the perturbative sense, gauge invariance gets restored also outside perturbation theory.

## 7 Triviality

While two dimensional chiral abelian gauge theories have a coupling dependent continuum limit<sup>4</sup>, four dimensional chiral abelian gauge theories are “trivial”

---

<sup>4</sup> Chiral two dimensional models, for compact  $U(1)$ , on a finite torus, have been studied in the overlap formalism in [9]. The formulation there is somewhat different from the one presented here, so that it be well adapted to numerical simulation. Using it, it was shown that such subtle phenomena as the appearance of massless fermionic composite particles, just so that 't Hooft's consistency relations be satisfied, indeed takes place for properly defined lattice models.

because the physical coupling is forced to vanish if one takes the ultraviolet cutoff to infinity. An anomaly free chiral  $U(1)$  theory, just like QED or the Higgs sector [10] of the minimal standard model, does not, strictly speaking, have an interacting continuum limit. Since the continuum limit is also jeopardized by lack of anomaly cancelation it is necessary to understand the difference between the two kinds of “trivialities” in four dimensions [2].

In four dimensions it is not necessary to deal with both left and right Weyl fermions, because a left handed fermion can be viewed as a right handed one in the conjugate representation. The anomaly cancelation condition can be written then as  $\sum_{\text{left handed fermions}} q^3 = 0$ ; if it is not satisfied, there is an obstruction to finding a section in the compounded bundle over  $A$  that descends into a section over gauge orbits [8]. This generalizes to the non-abelian case [11].

The difference between the two cases of triviality can be expressed by employing the effective field theory framework. Whether anomalies cancel or not, we can write down a theory which interacts but also has an ultraviolet cutoff  $\Lambda$ . This theory is adequate to describe to a given finite accuracy processes of energy up to some physical scale  $E_{\text{ph}}$ . Clearly,  $E_{\text{ph}} < \Lambda$ . For the accuracy of this description to be sensible the physical coupling  $e^{\text{ph}}$  cannot be too large. If anomalies do not cancel the restriction has the structure:

$$e^{\text{ph}} \leq c_1 \left( \frac{E^{\text{ph}}}{\Lambda} \right)^{\frac{1}{3}}, \quad (40)$$

where  $c_1$  is some constant. If anomalies do cancel the restriction is

$$e^{\text{ph}} \leq \frac{c_2}{\log \left( \frac{\Lambda}{E^{\text{ph}}} \right)}. \quad (41)$$

So, in both cases one cannot take  $\Lambda$  to infinity while keeping  $e^{\text{ph}}$  non-zero, but the restriction is much weaker if anomalies cancel.

## 8 Summary

There is little doubt left that indeed, as anticipated by most particle physics theorists, anomaly free, asymptotically free four dimensional chiral gauge theories have consistent continuum limits and exist outside perturbation theory. These are non-trivial theories, with no free parameters. These theories play an important role in our understanding of Nature, and what kind of mechanisms make them appear naturally at energies presently accessible is an important issue which may hold some hints about how things work in the real world.

## Acknowledgments

This research was supported in part by the DOE under grant # DE-FG05-96ER40559. I wish to thank the organizers of the “Quantization, Gauge Theory and Strings” dedicated to the memory of Professor Efim Fradkin for the invitation to participate, and for the hospitality extended.

## References

- [1] Proceedings of Chiral’99, The Workshop on Chiral Gauge Theories, *Chinese Journal of Physics*, Volume 38, Number 3, June 2000, Part II.
- [2] H. Neuberger, hep-lat/0002032.
- [3] H. Neuberger, *Phys. Rev. D* **61**, 085015 (2000).
- [4] H. Neuberger, *Phys. Lett. B* **417**, 141 (1998); *Phys. Rev. D* **57**, 5417 (1998); *Phys. Rev. Lett.* **81**, 4060 (1998).
- [5] R. Narayanan, H. Neuberger, *Phys. Lett. B* **302**, 62 (1993); *Nucl. Phys. B* **412**, 574 (1991); *Nucl. Phys. B* **443**, 305 (1995); *Phys. Rev. Lett.* **71**, 3251 (1993).
- [6] D. Kaplan *Phys. Lett. B* **288**, 342 (1992).
- [7] S. A. Frolov, A. A. Slavnov, *Phys. Lett. B* **303**, 344 (1993).
- [8] H. Neuberger, *Phys. Rev. D* **59**, 085006 (1999).
- [9] R. Narayanan, H. Neuberger, *Phys. Lett. B* **402**, 320 (1997).
- [10] H. Neuberger, *Phys. Rev. Lett.* **60**, 889 (1988);  
U. Heller, M. Klomfass, H. Neuberger, P. Vranas, *Nucl. Phys. B* **405**, 555 (1993);  
G. Bhanot, K. Bitar, U. Heller, H. Neuberger, *Nucl. Phys. B* **353**, 551 (1991).
- [11] D. Adams, hep-lat/0004015.